The American Academy of Ophthalmology is the world's largest association of eye physicians and surgeons. A global community of 32,000 medical doctors, we protect sight and empower lives by setting the standards for ophthalmic education and advocating for our patients and the public. For more information, visit www.aao.org.

COMPLIMENTS OF:

Get more information about macular hole from EyeSmart—provided by the American Academy of Ophthalmology—at aao.org/macular-hole-link.

©2022 American Academy of Ophthalmology 051198-9 Academy reviewed 09/22

978-1-61525-521-4

AMERICAN ACADEMY
OF OPHTHALMOLOGY®

- You may be asked to keep your face down in a specific position for a period of time, depending on your surgeon's preference. This is to keep the gas bubble in place to heal properly. Talk with your ophthalmologist about when you can do your everyday tasks again.
- You cannot fly in an airplane, go to mountains/high altitudes or scuba dive until the gas bubble is gone. This is because going up quickly in altitude can make eye pressure rise. That can cause problems with the bubble.
- If you need to have any other type of surgery, be sure to tell your doctor before surgery that you have a gas bubble in your eye.
- Your vision will improve as the macular hole closes. It may take several months for the hole to finish healing. How much vision you get back depends on the size of your macular hole. It also depends on how long the hole was there before you had surgery.

What are vitrectomy surgery risks?

Like any surgery, vitrectomy has risks. They include:

- Eye infection
- Bleeding in your eye
- A detached retina (where the retina lifts away from the back of the eye)
- Glaucoma, when pressure increases inside the eye
- Cataract, when the lens in your eye becomes cloudy


Your ophthalmologist will talk about these risks and how vitrectomy surgery may help you.

Summary

Macular hole is a circular opening in your macula, affecting your central vision. It is treated with a surgery called vitrectomy. Your eye surgeon removes the vitreous that is pulling on your macula.



Macular Hole



What is a macular hole?


Macular hole is when a circular opening forms in your **macula**. As the hole forms, things in your central vision will look blurry, wavy or distorted. As the hole grows, a dark or blind spot appears in your central vision. A macular hole does not affect your peripheral (side) vision.

Eye Words to Know

Retina: Layer of cells lining the back wall inside the eye. It senses light and sends signals to the brain so you can see.

Macula: Small but important area in the center of the retina. You need the macula to clearly see details and colors of objects in front of you.


Vitreous: Clear, gel-like substance that fills the inside of your eye. The vitreous helps the eye maintain its shape and also helps send light to the retina.




What causes a macular hole?

Age is the most common cause of macular hole. As you get older, the **vitreous** begins to shrink and pull away from the retina. Usually the vitreous pulls away with no problems. But sometimes the vitreous can stick to the **retina**. This causes the macula to stretch and a hole to form.

Sometimes a macular hole can form when the macula swells from other eye disease. Or it can be caused by an eye injury.





How is a macular hole diagnosed?

Your ophthalmologist will put drops in your eye to dilate (widen) your pupil. This allows them to look through a special lens at the inside of your eye.

Then they will take pictures of your eye using optical coherence tomography (OCT). With OCT, a machine scans the back of your eye. This provides very detailed pictures of the retina and macula. Your ophthalmologist studies these pictures to check for problems.

How is a macular hole treated?

Surgery called **vitrectomy** is the best way to treat a macular hole. Your ophthalmologist removes the vitreous that is pulling on your macula. Then they put a gas bubble inside the eye. This bubble helps flatten the macular hole and hold it in place while your eye heals. The gas bubble slowly goes away on its own, and is replaced by fluid your eye naturally produces.

Things to know about vitrectomy surgery for macular hole:

- Your eye may hurt after surgery. Your surgeon will have you take medicine to help with pain.
- You will need to wear an eye patch for a short time. You also will need to put drops in your eye.

2 3