



Uveitis

Swelling of the uvea toward the back of the eye is called **posterior uveitis**. Symptoms can develop gradually and last for many years.

In severe cases, all layers may be involved.

What causes uveitis?

Doctors do not always know what causes uveitis. You are more likely to get uveitis if you have or have had:

- infections such as shingles virus, herpes simplex virus, syphilis, Lyme disease, and parasites such as toxoplasmosis.
- a systemic inflammatory disease such as inflammatory bowel disease (IBD), rheumatoid arthritis or lupus
- an eye injury

Smoking (cigarettes, cigars or pipes) also increases your risk of getting uveitis.

How is uveitis diagnosed?

Your ophthalmologist will examine the inside of your eye. Since uveitis is often connected with other diseases or conditions, some tests may be needed. They may include a physical exam, blood or skin tests, examination of eye fluids, and imaging tests, such as X-rays. Your

ophthalmologist may ask about other diseases or health problems you have had.

How is uveitis treated?

Uveitis needs to be treated right away to prevent lasting problems.


Ophthalmologists often treat uveitis with eye drop medicine that reduces inflammation (corticosteroids). They may also use an eye drop to widen (dilate) the pupil, which helps reduce pain and swelling. Sometimes medicine may need to be given by injection (shots) or taken by mouth. In some cases, your ophthalmologist will work with a special kind of doctor called a rheumatologist to help treat you.

Summary

Uveitis is when the middle layer of the eyeball (uvea) gets inflamed. You are more likely to get uveitis if you have had certain diseases, viruses or an eye injury.


Symptoms include a red eye, pain and blurry vision. This is a serious condition that must be treated immediately to save vision. Treatment often includes eye drop medicine to reduce swelling and pain.

If you have any questions about your eyes or your vision, speak with your ophthalmologist. They are committed to protecting your sight.



What is uveitis?

Uveitis occurs when the middle layer of the eyeball gets inflamed (red and swollen). This layer, called the uvea, has many blood vessels that nourish the eye. Uveitis can damage vital eye tissue, leading to permanent vision loss.



What are uveitis symptoms?

Uveitis can develop suddenly. Symptoms can include:

- having a red eye with or without pain
- being very sensitive to bright light
- having blurry vision
- seeing “floaters” (specks or moving clouds in your vision) all of a sudden

Contact your ophthalmologist right away if you notice any of these symptoms.

Three types of uveitis

There are 3 types of uveitis. They are based on which part of the uvea is affected.

Swelling of the uvea near the front of the eye is called **anterior uveitis**. It starts suddenly and symptoms can last many weeks. Some forms of anterior uveitis are ongoing, while others go away but keep coming back.

Swelling of the uvea in the middle of the eye is called **intermediate uveitis**. Symptoms can last for a few weeks to many years. This form can go through cycles of getting better, then getting worse.

The American Academy of Ophthalmology is the world’s largest association of eye physicians and surgeons. A global community of 32,000 medical doctors, we protect sight and empower lives by setting the standards for ophthalmic education and advocating for our patients and the public. For more information, visit www.aao.org.

COMPLIMENTS OF:

Get more information about uveitis from EyeSmart—provided by the American Academy of Ophthalmology—at aao.org/uveitis-link.

©2023 American Academy of Ophthalmology

051219-10 Academy reviewed 03/23

978-1-61525-542-9



AMERICAN ACADEMY
OF OPHTHALMOLOGY®